

ACTOR OF A CROSSED MODULE OF LEIBNIZ ALGEBRAS

JOSÉ MANUEL CASAS, RAFAEL FERNÁNDEZ-CASADO,
XABIER GARCÍA-MARTÍNEZ, EMZAR KHMALADZE

ABSTRACT. We extend to the category of crossed modules of Leibniz algebras the notion of biderivation via the action of a Leibniz algebra. This results into a pair of Leibniz algebras which allow us to construct an object which is the actor under certain circumstances. Additionally, we give a description of an action in the category of crossed modules of Leibniz algebras in terms of equations. Finally, we check that, under the aforementioned conditions, the kernel of the canonical map from a crossed module to its actor coincides with the center and we introduce the notions of crossed module of inner and outer biderivations.

1. Introduction

Let H be a group and $\text{Aut}(H)$ be the group of automorphisms. For every action of a group G on H there is a unique group homomorphism $\beta: G \rightarrow \text{Aut}(H)$ with ${}^g h = \beta(g)(h)$ for all $g \in G$ and $h \in H$. Conversely, every group homomorphism from G to $\text{Aut}(H)$ induces an action of G on H . Therefore it is equivalent to consider group actions of G on H or group homomorphisms from G to $\text{Aut}(H)$. By this fact, the group of automorphisms is called the *actor* in the category of groups. Its analogue in the category of Lie algebras is the Lie algebra of derivations.

More generally, for any semi-abelian category \mathbf{C} , the existence of the actor is determined by the representability of the contravariant functor from \mathbf{C} to the category of sets, sending an object Y to the set of actions of Y on X [2]. That is, the existence of an object $\text{Act}(X)$ such that the set of actions of Y on X is isomorphic to $\text{Hom}_{\mathbf{C}}(Y, \text{Act}(X))$. If this object exists for every X in \mathbf{C} , the category is said to be *action representable* [3], and the object $\text{Act}(X)$ is called the *actor* [15] (also known as *split extension classifier* [3]).

Groups and Lie algebras are examples of categories of interest, introduced by Orzech in [16]. For these categories (see [14] for more examples), Casas, Datuashvili and Ladra [6] gave a procedure to construct an object that, under certain circumstances, plays the

The authors were supported by Ministerio de Economía y Competitividad (Spain), grant MTM2016-79661-P (European FEDER support included). The third author was also supported by an FPU scholarship, Ministerio de Educación, Cultura y Deporte (Spain). The fourth author was supported by Shota Rustaveli National Science Foundation, grant FR/189/5-113/14.

Received by the editors 2016-10-02 and, in final form, 2017-10-29.

Transmitted by Ieke Moerdijk. Published on 2018-01-04.

2010 Mathematics Subject Classification: 17A30, 17A32, 18A05, 18D05.

Key words and phrases: Leibniz algebra, crossed module, representation, actor.

© José Manuel Casas, Rafael Fernández-Casado,

Xabier García-Martínez, Emzar Khmaladze, 2018. Permission to copy for private use granted.

role of actor. For the particular case of Leibniz algebras (resp. associative algebras) that object is the Leibniz algebra of biderivations (resp. the algebra of bimultipliers).

In [15], Norrie extended the definition of actor to the 2-dimensional case by giving a description of the corresponding object in the category of crossed modules of groups. The analogue construction for the category of crossed modules of Lie algebras is given in [10]. Regarding the category of crossed modules of Leibniz algebras, it is not a category of interest, but it is equivalent to the category of cat^1 -Leibniz algebras (see for example [9]), which is itself a modified category of interest in the sense of [4]. Therefore it makes sense to study representability of actions in such category under the context of modified categories of interest, as it is done in [4] for crossed modules of associative algebras.

Bearing in mind the ease of the generalization of the actor in the category of groups and Lie algebras to crossed modules, together with the role of the Leibniz algebra of biderivations, it makes sense to assume that the analogous object in the category of crossed modules of Leibniz algebras will be the actor only under certain hypotheses. In [8] the authors gave an equivalent description of an action of a crossed module of groups in terms of equations. A similar description is done for an action of a crossed module of Lie algebras (see [5]). In order to extend the notion of actor to crossed modules of Leibniz algebras, we generalize the concept of biderivation to the 2-dimensional case, describe an action in that category in terms of equations and give sufficient conditions for the described object to be the actor.

The article is organized as follows: In Section 2 we recall some basic definitions on actions and crossed modules of Leibniz algebras. In Section 3 we construct an object that extends the Leibniz algebra of biderivations to the category of crossed modules of Leibniz algebras (Theorem 3.9) and give a description of an action in such category in terms of equations. In Section 4 we find sufficient conditions for the previous object to be the actor of a given crossed module of Leibniz algebras (Theorem 4.3). Finally, in Section 5 we prove that the kernel of the canonical homomorphism from a crossed module of Leibniz algebras to its actor coincides with the center of the given crossed module. Additionally, we introduce the notions of crossed module of inner and outer biderivations and show that, given a short exact sequence in the category of crossed modules of Leibniz algebras, it can be extended to a commutative diagram including the actor and the inner and outer biderivations.

2. Preliminaries

In this section we recall some needed basic definitions. Throughout the paper we fix a commutative ring with unit \mathbf{k} . All algebras are considered over \mathbf{k} .

2.1. DEFINITION. [12] *A Leibniz algebra \mathfrak{p} is a \mathbf{k} -module together with a bilinear operation $[\ , \]: \mathfrak{p} \times \mathfrak{p} \rightarrow \mathfrak{p}$, called the Leibniz bracket, which satisfies the Leibniz identity:*

$$[[p_1, p_2], p_3] = [p_1, [p_2, p_3]] + [[p_1, p_3], p_2],$$

for all $p_1, p_2, p_3 \in \mathfrak{p}$.

A homomorphism of Leibniz algebras is a \mathbf{k} -linear map that preserves the bracket.

We denote by $\text{Ann}(\mathfrak{p})$ (resp. $[\mathfrak{p}, \mathfrak{p}]$) the *annihilator* (resp. commutator) of \mathfrak{p} , that is the subspace of \mathfrak{p} generated by

$$\begin{aligned} & \{p_1 \in \mathfrak{p} \mid [p_1, p_2] = [p_2, p_1] = 0, \text{ for all } p_2 \in \mathfrak{p}\} \\ & \text{(resp. } \{[p_1, p_2] \mid \text{for all } p_1, p_2 \in \mathfrak{p}\}) \end{aligned}$$

It is obvious that both $\text{Ann}(\mathfrak{p})$ and $[\mathfrak{p}, \mathfrak{p}]$ are ideals of \mathfrak{p} .

2.2. DEFINITION. [13] *Let \mathfrak{p} and \mathfrak{m} be two Leibniz algebras. An action of \mathfrak{p} on \mathfrak{m} consists of a pair of bilinear maps, $\mathfrak{p} \times \mathfrak{m} \rightarrow \mathfrak{m}$, $(p, m) \mapsto {}^p m$ and $\mathfrak{m} \times \mathfrak{p} \rightarrow \mathfrak{m}$, $(m, p) \mapsto m^p$, such that*

$$\begin{aligned} {}^p[m, m'] &= [{}^p m, m'] - [{}^p m', m], \\ [m, {}^p m'] &= [m^p, m'] - [m, m']^p, \\ [m, m'^p] &= [m, m']^p - [m^p, m'], \\ m^{[p, p']} &= (m^p)^{p'} - (m^{p'})^p, \\ {}^p(m^{p'}) &= ({}^p m)^{p'} - [{}^p, p']m, \\ {}^p(p' m) &= [{}^p, p']m - ({}^p m)^{p'}, \end{aligned}$$

for all $m, m' \in \mathfrak{m}$ and $p, p' \in \mathfrak{p}$.

Given an action of a Leibniz algebra \mathfrak{p} on \mathfrak{m} , we can consider the *semidirect product* Leibniz algebra $\mathfrak{m} \rtimes \mathfrak{p}$, which consists of the \mathbf{k} -module $\mathfrak{m} \oplus \mathfrak{p}$ together with the Leibniz bracket given by

$$[(m, p), (m', p')] = ([m, m'] + {}^p m' + m^{p'}, [p, p']),$$

for all $(m, p), (m', p') \in \mathfrak{m} \oplus \mathfrak{p}$.

2.3. DEFINITION. [13] *A crossed module of Leibniz algebras (or Leibniz crossed module, for short) $(\mathfrak{m}, \mathfrak{p}, \eta)$ is a homomorphism of Leibniz algebras $\eta: \mathfrak{m} \rightarrow \mathfrak{p}$ together with an action of \mathfrak{p} on \mathfrak{m} such that*

$$\eta({}^p m) = [p, \eta(m)] \quad \text{and} \quad \eta(m^p) = [\eta(m), p], \quad (\text{XLb1})$$

$$\eta^{(m)} m' = [m, m'] = m^{\eta(m')}, \quad (\text{XLb2})$$

for all $m, m' \in \mathfrak{m}$, $p \in \mathfrak{p}$.

A homomorphism of Leibniz crossed modules (φ, ψ) from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $(\mathfrak{n}, \mathfrak{q}, \mu)$ is a pair of Leibniz homomorphisms, $\varphi: \mathfrak{m} \rightarrow \mathfrak{n}$ and $\psi: \mathfrak{p} \rightarrow \mathfrak{q}$, such that they commute with η and μ and they respect the actions, that is $\varphi({}^p m) = {}^{\psi(p)} \varphi(m)$ and $\varphi(m^p) = \varphi(m)^{\psi(p)}$ for all $m \in \mathfrak{m}$, $p \in \mathfrak{p}$.

Identity (XLb1) will be called *equivariance* and (XLb2) *Peiffer identity*. We will denote by **XLb** the category of Leibniz crossed modules and homomorphisms of Leibniz crossed modules.

Since our aim is to construct a 2-dimensional generalization of the actor in the category of Leibniz algebras, let us first recall the following definitions.

2.4. DEFINITION. [12] *Let \mathfrak{m} be a Leibniz algebra. A biderivation of \mathfrak{m} is a pair (d, D) of \mathbf{k} -linear maps $d, D: \mathfrak{m} \rightarrow \mathfrak{m}$ such that*

$$d([m, m']) = [d(m), m'] + [m, d(m')], \quad (1)$$

$$D([m, m']) = [D(m), m'] - [D(m'), m], \quad (2)$$

$$[m, d(m')] = [m, D(m')], \quad (3)$$

for all $m, m' \in \mathfrak{m}$.

We will denote by $\text{Bider}(\mathfrak{m})$ the set of all biderivations of \mathfrak{m} . It is a Leibniz algebra with the obvious \mathbf{k} -module structure and the Leibniz bracket given by

$$[(d_1, D_1), (d_2, D_2)] = (d_1 d_2 - d_2 d_1, D_1 d_2 - d_2 D_1).$$

It is not difficult to check that, given an element $m \in \mathfrak{m}$, the pair $(\text{ad}(m), \text{Ad}(m))$, with $\text{ad}(m)(m') = -[m', m]$ and $\text{Ad}(m)(m') = [m, m']$ for all $m' \in \mathfrak{m}$, is a biderivation. The pair $(\text{ad}(m), \text{Ad}(m))$ is called *inner biderivation* of m .

3. The main construction

In this section we extend to crossed modules the Leibniz algebra of biderivations. First we need to translate the notion of a biderivation of a Leibniz algebra into a biderivation between two Leibniz algebras via the action.

3.1. DEFINITION. *Given an action of Leibniz algebras of \mathfrak{q} on \mathfrak{n} , the set of biderivations from \mathfrak{q} to \mathfrak{n} , denoted by $\text{Bider}(\mathfrak{q}, \mathfrak{n})$, consists of all the pairs (d, D) of \mathbf{k} -linear maps, $d, D: \mathfrak{q} \rightarrow \mathfrak{n}$, such that*

$$d([q, q']) = d(q)^{q'} + {}^q d(q'), \quad (4)$$

$$D([q, q']) = D(q)^{q'} - D(q')^q, \quad (5)$$

$${}^q d(q') = {}^q D(q'), \quad (6)$$

for all $q, q' \in \mathfrak{q}$.

Given $n \in \mathfrak{n}$, the pair of \mathbf{k} -linear maps $(\text{ad}(n), \text{Ad}(n))$, where $\text{ad}(n)(q) = -{}^q n$ and $\text{Ad}(n)(q) = n^q$ for all $q \in \mathfrak{q}$, is clearly a biderivation from \mathfrak{q} to \mathfrak{n} . Observe that $\text{Bider}(\mathfrak{q}, \mathfrak{q})$, with the action of \mathfrak{q} on itself defined by its Leibniz bracket, is exactly $\text{Bider}(\mathfrak{q})$.

Let us assume for the rest of the article that $(\mathfrak{n}, \mathfrak{q}, \mu)$ is a Leibniz crossed module. One can easily check the following result.

3.2. LEMMA. *Let $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. Then $(d\mu, D\mu) \in \text{Bider}(\mathfrak{n})$ and $(\mu d, \mu D) \in \text{Bider}(\mathfrak{q})$.*

We also have the following result.

3.3. LEMMA. *Let $(d_1, D_1), (d_2, D_2) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. Then*

$$\begin{aligned} (D_1\mu d_2(q))^{q'} &= (D_1\mu D_2(q))^{q'}, \\ {}^q(D_1\mu d_2(q')) &= {}^q(D_1\mu D_2(q')), \end{aligned}$$

for all $q, q' \in \mathfrak{q}$.

PROOF. Let $q, q' \in \mathfrak{q}$ and $(d_1, D_1), (d_2, D_2) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. According to the identity (6) for (d_2, D_2) , ${}^{q'}d_2(q) = {}^{q'}D_2(q)$, so $D_1\mu({}^{q'}d_2(q)) = D_1\mu({}^{q'}D_2(q))$. Due to (5) and the equivariance of $(\mathfrak{q}, \mathfrak{n}, \mu)$, one can easily derive that

$$D_1(q')^{\mu d_2(q)} - (D_1\mu d_2(q))^{q'} = D_1(q')^{\mu D_2(q)} - (D_1\mu D_2(q))^{q'}.$$

By the Peiffer identity and (6) for (d_2, D_2) , $D_1(q')^{\mu d_2(q)} = D_1(q')^{\mu D_2(q)}$. Then $(D_1\mu d_2(q))^{q'} = (D_1\mu D_2(q))^{q'}$.

The other identity can be proved similarly by using (4) and (6). ■

$\text{Bider}(\mathfrak{q}, \mathfrak{n})$ has an obvious \mathbf{k} -module structure. Regarding its Leibniz structure, it is described in the next proposition.

3.4. PROPOSITION. *$\text{Bider}(\mathfrak{q}, \mathfrak{n})$ is a Leibniz algebra with the bracket given by*

$$[(d_1, D_1), (d_2, D_2)] = (d_1\mu d_2 - d_2\mu d_1, D_1\mu d_2 - d_2\mu D_1) \quad (7)$$

for all $(d_1, D_1), (d_2, D_2) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$.

PROOF. It follows directly from Lemma 3.3. ■

Now we state the following definition.

3.5. DEFINITION. *The set of biderivations of the Leibniz crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$, denoted by $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$, consists of all quadruples $((\sigma_1, \theta_1), (\sigma_2, \theta_2))$ such that*

$$(\sigma_1, \theta_1) \in \text{Bider}(\mathfrak{n}) \quad \text{and} \quad (\sigma_2, \theta_2) \in \text{Bider}(\mathfrak{q}), \quad (8)$$

$$\mu\sigma_1 = \sigma_2\mu \quad \text{and} \quad \mu\theta_1 = \theta_2\mu, \quad (9)$$

$$\sigma_1({}^q n) = \sigma_2({}^q n) + {}^q\sigma_1(n), \quad (10)$$

$$\sigma_1(n^q) = \sigma_1(n)^q + n^{\sigma_2(q)}, \quad (11)$$

$$\theta_1({}^q n) = \theta_2({}^q n) - \theta_1(n)^q, \quad (12)$$

$$\theta_1(n^q) = \theta_1(n)^q - \theta_2(q)n, \quad (13)$$

$${}^q\sigma_1(n) = {}^q\theta_1(n), \quad (14)$$

$$n^{\sigma_2(q)} = n^{\theta_2(q)}, \quad (15)$$

for all $n \in \mathfrak{n}, q \in \mathfrak{q}$.

Given $q \in \mathfrak{q}$, it can be readily checked that $((\sigma_1^q, \theta_1^q), (\sigma_2^q, \theta_2^q))$, where

$$\begin{aligned} \sigma_{1,q}(n) &= -n^q, & \theta_{1,q}(n) &= {}^q n, \\ \sigma_{2,q}(q') &= -[q', q], & \theta_{2,q}(q') &= [q, q'], \end{aligned}$$

is a biderivation of the crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$.

The following lemma is necessary in order to prove that $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ is indeed a Leibniz algebra.

3.6. LEMMA. *Let $((\sigma_1, \theta_1), (\sigma_2, \theta_2)), ((\sigma'_1, \theta'_1), (\sigma'_2, \theta'_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$, $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. Then*

$$\begin{aligned} (D\sigma_2(q))^{q'} &= (D\theta_2(q))^{q'}, & [D\sigma_2(q), n] &= [D\theta_2(q), n], & (\theta_1\sigma'_1(n))^q &= (\theta_1\theta'_1(n))^q, \\ {}^q(D\sigma_2(q')) &= {}^q(D\theta_2(q')), & [n, D\sigma_2(q)] &= [n, D\theta_2(q)], & {}^q(\theta_1\sigma'_1(n)) &= {}^q(\theta_1\theta'_1(n)), \\ (\theta_1d(q))^{q'} &= (\theta_1D(q))^{q'}, & [\theta_1d(q), n] &= [\theta_1D(q), n], & \theta_2\sigma'_2(q)_n &= \theta_2\theta'_2(q)_n, \\ {}^q(\theta_1d(q')) &= {}^q(\theta_1D(q')), & [n, \theta_1d(q)] &= [n, \theta_1D(q)], & n^{\theta_2\sigma'_2(q)} &= n^{\theta_2\theta'_2(q)}, \end{aligned}$$

for all $n \in \mathfrak{n}$, $q, q' \in \mathfrak{q}$.

PROOF. Let us show how to prove the first identity; the rest of them can be checked similarly. Let $q, q' \in \mathfrak{q}$, $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$ and $((\sigma_1, \theta_1), (\sigma_2, \theta_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$. Since (σ_2, θ_2) is a biderivation of \mathfrak{q} , we have that $[q', \sigma_2(q)] = [q', \theta_2(q)]$. Therefore $D([q', \sigma_2(q)]) = D([q', \theta_2(q)])$. Directly from (5), we get that

$$D(q')^{\sigma_2(q)} - (D\sigma_2(q))^{q'} = D(q')^{\theta_2(q)} - (D\theta_2(q))^{q'}.$$

Thus, due to (15), $D(q')^{\sigma_2(q)} = D(q')^{\theta_2(q)}$. Hence, $(D\sigma_2(q))^{q'} = (D\theta_2(q))^{q'}$. ■

The \mathbf{k} -module structure of $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ is evident, while its Leibniz structure is described as follows.

3.7. PROPOSITION. *$\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ is a Leibniz algebra with the bracket given by*

$$\begin{aligned} [((\sigma_1, \theta_1), (\sigma_2, \theta_2)), ((\sigma'_1, \theta'_1), (\sigma'_2, \theta'_2))] &= ([(\sigma_1, \theta_1), (\sigma'_1, \theta'_1)], [(\sigma_2, \theta_2), (\sigma'_2, \theta'_2)]) \\ &= ((\sigma_1\sigma'_1 - \sigma'_1\sigma_1, \theta_1\sigma'_1 - \sigma'_1\theta_1), (\sigma_2\sigma'_2 - \sigma'_2\sigma_2, \theta_2\sigma'_2 - \sigma'_2\theta_2)), \end{aligned} \quad (16)$$

for all $((\sigma_1, \theta_1), (\sigma_2, \theta_2)), ((\sigma'_1, \theta'_1), (\sigma'_2, \theta'_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$.

PROOF. It follows directly from Lemma 3.6. ■

3.8. PROPOSITION. *The \mathbf{k} -linear map $\Delta: \text{Bider}(\mathfrak{q}, \mathfrak{n}) \rightarrow \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$, given by $(d, D) \mapsto ((d\mu, D\mu), (\mu d, \mu D))$ is a homomorphism of Leibniz algebras.*

PROOF. Δ is well defined due to Lemma 3.2, while checking that it is a homomorphism of Leibniz algebras is a matter of straightforward calculations. ■

Since we aspire to make Δ into a Leibniz crossed module, we need to define an action of $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ on $\text{Bider}(\mathfrak{q}, \mathfrak{n})$.

3.9. THEOREM. *There is an action of $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ on $\text{Bider}(\mathfrak{q}, \mathfrak{n})$ given by:*

$$((\sigma_1, \theta_1), (\sigma_2, \theta_2))(d, D) = (\sigma_1 d - d\sigma_2, \theta_1 d - d\theta_2), \quad (17)$$

$$(d, D)^{((\sigma_1, \theta_1), (\sigma_2, \theta_2))} = (d\sigma_2 - \sigma_1 d, D\sigma_2 - \sigma_1 D), \quad (18)$$

for all $((\sigma_1, \theta_1), (\sigma_2, \theta_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$, $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. Moreover, the Leibniz homomorphism Δ (see Proposition 3.8) together with the above action is a Leibniz crossed module.

PROOF. Let $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$ and $((\sigma_1, \theta_1), (\sigma_2, \theta_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$. Checking that both $(\sigma_1 d - d\sigma_2, \theta_1 d - d\theta_2)$ and $(d\sigma_2 - \sigma_1 d, D\sigma_2 - \sigma_1 D)$ satisfy conditions (4) and (5) requires the combined use of the properties satisfied by the elements in $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ and (d, D) , but calculations are fairly straightforward. As an example, we show how to prove that $(\sigma_1 d - d\sigma_2, \theta_1 d - d\theta_2)$ verifies (4). Let $q, q' \in \mathfrak{q}$. Then

$$\begin{aligned} (\sigma_1 d - d\sigma_2)([q, q']) &= \sigma_1(d(q)^{q'} + {}^q d(q')) - d([\sigma_2(q), q'] + [q, \sigma_2(q')]) \\ &= (\sigma_1 d(q))^{q'} + d(q)^{\sigma_2(q')} + \sigma_2(q)d(q') + {}^q(\sigma_1 d(q')) \\ &\quad - (d\sigma_2(q))^{q'} - \sigma_2(q)d(q') - d(q)^{\sigma_2(q')} - {}^q(d\sigma_2(q')) \\ &= ((\sigma_1 d - d\sigma_2)(q))^{q'} + {}^q((\sigma_1 d - d\sigma_2)(q')). \end{aligned}$$

As for condition (6), in the case of $(\sigma_1 d - d\sigma_2, \theta_1 d - d\theta_2)$, it follows from (14), the identity (6) for (d, D) and the second identity in the first column from Lemma 3.6. Namely,

$$\begin{aligned} {}^q((\sigma_1 d - d\sigma_2)(q')) &= {}^q(\sigma_1 d(q')) - {}^q(d\sigma_2(q')) = {}^q(\theta_1 d(q')) - {}^q(D\sigma_2(q')) \\ &= {}^q(\theta_1 d(q')) - {}^q(D\theta_2(q')) = {}^q(\theta_1 d(q')) - {}^q(d\theta_2(q')), \end{aligned}$$

for all $q, q' \in \mathfrak{q}$. A similar procedure allows to prove that $(d\sigma_2 - \sigma_1 d, D\sigma_2 - \sigma_1 D)$ satisfies condition (6) as well.

Routine calculations show that (17) and (18) together with the definition of the brackets in $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ and $\text{Bider}(\mathfrak{q}, \mathfrak{n})$ provide an action of Leibniz algebras.

It only remains to prove that Δ satisfies the equivariance and the Peiffer identity. It is immediate to check that

$$\Delta^{((\sigma_1, \theta_1), (\sigma_2, \theta_2))}(d, D) = ((\sigma_1 d\mu - d\sigma_2\mu, \theta_1 d\mu - d\theta_2\mu), (\mu\sigma_1 d - \mu d\sigma_2, \mu\theta_1 d - \mu d\theta_2)), \quad (19)$$

while

$$\begin{aligned} [((\sigma_1, \theta_1), (\sigma_2, \theta_2)), \Delta(d, D)] &= ((\sigma_1 d\mu - d\mu\sigma_1, \theta_1 d\mu - d\mu\theta_1), \\ &\quad (\sigma_2\mu d - \mu d\sigma_2, \theta_2\mu d - \mu d\theta_2)). \end{aligned} \quad (20)$$

Condition (9) guarantees that (19) = (20). The other identity can be checked similarly. The Peiffer identity follows immediately from (17) and (18) along the definition of Δ and the bracket in $\text{Bider}(\mathfrak{q}, \mathfrak{n})$. \blacksquare

4. The actor

In [16], Orzech introduced the notion of category of interest, which is nothing but a category of groups with operations verifying two extra conditions. \mathbf{Lb} is a category of interest, although \mathbf{XLb} is not. Nevertheless, it is equivalent to the category of cat^1 -Leibniz algebras (see for example [9]), which is itself a modified category of interest in the sense of [4]. So it makes sense to study representability of actions in \mathbf{XLb} under the context of modified categories of interest, as it is done in [4] for crossed modules of associative algebras. However, since \mathbf{XLb} is an example of semi-abelian category, and an action is the same as a split extension in any semi-abelian category [2, Lemma 1.3], we choose a different, more combinatorial approach to the problem, by constructing the semidirect product (split extension) of Leibniz crossed modules.

As we mention in the introduction, we use the term *actor* (as in [4, 6]) for an object which represents actions in a semi-abelian category, the general definition of which is known from [3] under the name *split extension classifier*.

We need to remark that, given a Leibniz algebra \mathfrak{m} , $\text{Bider}(\mathfrak{m})$ is the actor of \mathfrak{m} under certain conditions. In particular, the following result is proved in [6].

4.1. PROPOSITION. [6] *Let \mathfrak{m} be a Leibniz algebra such that $\text{Ann}(\mathfrak{m}) = 0$ or $[\mathfrak{m}, \mathfrak{m}] = \mathfrak{m}$. Then $\text{Bider}(\mathfrak{m})$ is the actor of \mathfrak{m} .*

Bearing in mind the ease of the generalization of the actor in the category of groups and Lie algebras to crossed modules, together with the role of $\text{Bider}(\mathfrak{m})$ in regard to any Leibniz algebra \mathfrak{m} , it makes sense to consider $(\text{Bider}(\mathfrak{q}, \mathfrak{n}), \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu), \Delta)$ as a candidate for actor in \mathbf{XLb} , at least under certain conditions (see Proposition 4.1). However, it would be reckless to define an action of a Leibniz crossed module $(\mathfrak{m}, \mathfrak{p}, \eta)$ on $(\mathfrak{n}, \mathfrak{q}, \mu)$ as a homomorphism from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to the Leibniz crossed module $(\text{Bider}(\mathfrak{q}, \mathfrak{n}), \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu), \Delta)$, since we cannot ensure that the mentioned homomorphism induces a set of actions of $(\mathfrak{m}, \mathfrak{p}, \eta)$ on $(\mathfrak{n}, \mathfrak{q}, \mu)$ from which we can construct the semidirect product.

In [8, Proposition 2.1] the authors give an equivalent description of an action of a crossed module of groups in terms of equations. A similar description can be done for an action of a crossed module of Lie algebras (see [5]). This determines our approach to the problem. We consider a homomorphism from a Leibniz crossed module $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $(\text{Bider}(\mathfrak{q}, \mathfrak{n}), \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu), \Delta)$, which will be denoted by $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ from now on, and unravel all the properties satisfied by the mentioned homomorphism, transforming them into a set of equations. Then we check that the existence of that set of equations is equivalent to the existence of a homomorphism of Leibniz crossed modules from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ only under certain conditions. Finally we prove that those equations indeed describe a comprehensive set of actions by constructing the associated semidirect product, which is an object in \mathbf{XLb} .

4.2. LEMMA.

- (i) *Let \mathfrak{q} be a Leibniz algebra and $(\sigma, \theta), (\sigma', \theta') \in \text{Bider}(\mathfrak{q})$. If $\text{Ann}(\mathfrak{q}) = 0$ or $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$,*

then

$$\theta\sigma'(q) = \theta\theta'(q), \quad (21)$$

for all $q \in \mathfrak{q}$.

- (ii) Let $(\mathfrak{n}, \mathfrak{q}, \mu)$ be a Leibniz crossed module, $((\sigma_1, \theta_1), (\sigma_2, \theta_2)) \in \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ and $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$. If $\text{Ann}(\mathfrak{n}) = 0$ or $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$, then

$$D\sigma_2(q) = D\theta_2(q), \quad (22)$$

$$\theta_1 d(q) = \theta_1 D(q), \quad (23)$$

for all $q \in \mathfrak{q}$.

PROOF. Calculations in order to prove (i) are straightforward. Regarding (ii), $D\sigma_2(q) - D\theta_2(q)$ and $\theta_1 d(q) - \theta_1 D(q)$ are elements in $\text{Ann}(\mathfrak{n})$, immediately from the identities in the second column from Lemma 3.6. Therefore, if $\text{Ann}(\mathfrak{n}) = 0$, it is clear that (22) and (23) hold.

Let us now assume that $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$. Given $q, q' \in \mathfrak{q}$, directly from the fact that $(\sigma_2, \theta_2) \in \text{Bider}(\mathfrak{q})$ and $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$, we get that

$$D\theta_2([q, q']) = (D\theta_2(q))^{q'} - D(q')^{\theta_2(q)} - (D\theta_2(q'))^q + D(q)^{\theta_2(q')},$$

$$D\sigma_2([q, q']) = (D\sigma_2(q))^{q'} - D(q')^{\sigma_2(q)} + D(q)^{\sigma_2(q')} - (D\sigma_2(q'))^q.$$

Due to (15) and the first identity in the first column from Lemma 3.6, $D\theta_2([q, q']) = D\sigma_2([q, q'])$. By hypothesis, every element in \mathfrak{q} can be expressed as a linear combination of elements of the form $[q, q']$. This fact together with the linearity of D , σ_2 and θ_2 , guarantees that $D\theta_2(q) = D\sigma_2(q)$ for all $q \in \mathfrak{q}$. The identity (23) can be checked similarly by making use of (6), (12), (13) and the third identity in the first column from Lemma 3.6. ■

4.3. THEOREM. *Let $(\mathfrak{m}, \mathfrak{p}, \eta)$ and $(\mathfrak{n}, \mathfrak{q}, \mu)$ in \mathbf{XLb} . There exists a homomorphism of crossed modules from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $(\text{Bider}(\mathfrak{q}, \mathfrak{n}), \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu), \Delta)$, if the following conditions hold:*

- (i) *There are actions of the Leibniz algebra \mathfrak{p} (and so \mathfrak{m}) on the Leibniz algebras \mathfrak{n} and \mathfrak{q} . The homomorphism μ is \mathfrak{p} -equivariant, that is*

$$\mu({}^p n) = {}^p \mu(n), \quad (\text{LbEQ1})$$

$$\mu(n^p) = \mu(n)^p, \quad (\text{LbEQ2})$$

and the actions of \mathfrak{p} and \mathfrak{q} on \mathfrak{n} are compatible, that is

$$n^{(pq)} = (n^p)^q - (n^q)^p, \quad (\text{LbCOM1})$$

$${}^p(n^q) = ({}^p n)^q - ({}^p q)n, \quad (\text{LbCOM2})$$

$${}^p(qn) = ({}^p q)n - ({}^p n)^q, \quad (\text{LbCOM3})$$

$$n^{(qp)} = (n^q)^p - (n^p)^q, \quad (\text{LbCOM4})$$

$${}^q(n^p) = ({}^q n)^p - ({}^q p)n, \quad (\text{LbCOM5})$$

$${}^q(pn) = ({}^q p)n - ({}^q n)^p, \quad (\text{LbCOM6})$$

for all $n \in \mathfrak{n}$, $p \in \mathfrak{p}$ and $q \in \mathfrak{q}$.

(ii) There are two \mathbf{k} -bilinear maps $\xi_1: \mathfrak{m} \times \mathfrak{q} \rightarrow \mathfrak{n}$ and $\xi_2: \mathfrak{q} \times \mathfrak{m} \rightarrow \mathfrak{n}$ such that

$$\mu\xi_2(q, m) = q^m, \quad (\text{LbM1a})$$

$$\mu\xi_1(m, q) = {}^m q, \quad (\text{LbM1b})$$

$$\xi_2(\mu(n), m) = n^m, \quad (\text{LbM2a})$$

$$\xi_1(m, \mu(n)) = {}^m n, \quad (\text{LbM2b})$$

$$\xi_2(q, {}^p m) = \xi_2(q^p, m) - \xi_2(q, m)^p, \quad (\text{LbM3a})$$

$$\xi_1({}^p m, q) = \xi_2({}^p q, m) - {}^p \xi_2(q, m), \quad (\text{LbM3b})$$

$$\xi_2(q, m^p) = \xi_2(q, m)^p - \xi_2(q^p, m), \quad (\text{LbM3c})$$

$$\xi_1(m^p, q) = \xi_1(m, q)^p - \xi_1(m, q^p), \quad (\text{LbM3d})$$

$$\xi_2(q, [m, m']) = \xi_2(q, m)^{m'} - \xi_2(q, m')^m, \quad (\text{LbM4a})$$

$$\xi_1([m, m'], q) = \xi_1(m, q)^{m'} - {}^m \xi_2(q, m'), \quad (\text{LbM4b})$$

$$\xi_2([q, q'], m) = \xi_2(q, m)^{q'} + {}^q \xi_2(q', m), \quad (\text{LbM5a})$$

$$\xi_1(m, [q, q']) = \xi_1(m, q)^{q'} - \xi_1(m, q')^q, \quad (\text{LbM5b})$$

$${}^q \xi_1(m, q') = -{}^q \xi_2(q', m), \quad (\text{LbM5c})$$

$$\xi_1(m, {}^p q) = -\xi_1(m, q^p), \quad (\text{LbM6a})$$

$${}^p \xi_1(m, q) = -{}^p \xi_2(q, m), \quad (\text{LbM6b})$$

for all $m, m' \in \mathfrak{m}$, $n \in \mathfrak{n}$, $p \in \mathfrak{p}$, $q, q' \in \mathfrak{q}$.

Additionally, the converse statement is also true if one of the following conditions holds:

$$\text{Ann}(\mathfrak{n}) = 0 = \text{Ann}(\mathfrak{q}), \quad (\text{CON1})$$

$$\text{Ann}(\mathfrak{n}) = 0 \quad \text{and} \quad [\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}, \quad (\text{CON2})$$

$$[\mathfrak{n}, \mathfrak{n}] = \mathfrak{n} \quad \text{and} \quad [\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}. \quad (\text{CON3})$$

PROOF. Let us suppose that (i) and (ii) hold. It is possible to define a homomorphism of crossed modules (φ, ψ) from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ as follows. Given $m \in \mathfrak{m}$, $\varphi(m) = (d_m, D_m)$, with

$$d_m(q) = -\xi_2(q, m), \quad D_m(q) = \xi_1(m, q),$$

for all $q \in \mathfrak{q}$. On the other hand, for any $p \in \mathfrak{p}$, $\psi(p) = ((\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p}))$, with

$$\begin{aligned} \sigma_{1,p}(n) &= -n^p, & \theta_{1,p}(n) &= {}^p n, \\ \sigma_{2,p}(q) &= -q^p, & \theta_{2,p}(q) &= {}^p q, \end{aligned}$$

for all $n \in \mathfrak{n}$, $q \in \mathfrak{q}$. It follows directly from (LbM5a)–(LbM5c) that $(d_m, D_m) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$ for all $m \in \mathfrak{m}$. Besides, φ is clearly \mathbf{k} -linear and given $m, m' \in \mathfrak{m}$,

$$[\varphi(m), \varphi(m')] = [(d_m, D_m), (d_{m'}, D_{m'})] = [d_m \mu d_{m'} - d_{m'} \mu d_m, D_m \mu d_{m'} - d_{m'} \mu D_m].$$

For any $q \in \mathfrak{q}$,

$$\begin{aligned} d_m \mu d_{m'}(q) - d_{m'} \mu d_m(q) &= -\xi_2(\mu d_{m'}(q), m) + \xi_2(\mu d_m(q), m') \\ &= -d_{m'}(q)^m + d_m(q)^{m'} \\ &= \xi_2(q, m')^m - \xi_2(q, m)^{m'} \\ &= -\xi_2(q, [m, m']) = d_{[m, m']}(q), \end{aligned}$$

due to (LbM2a) and (LbM4a). Analogously, it can be easily checked the identity $(D_m \mu d_{m'} - d_{m'} \mu D_m)(q) = D_{[m, m']}(q)$ by making use of (LbM2a), (LbM2b) and (LbM4b). Hence, φ is a homomorphism of Leibniz algebras.

As for ψ , it is necessary to prove that $((\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p}))$ satisfies all the axioms from Definition 3.5 for any $p \in \mathfrak{p}$. The fact that $(\sigma_{1,p}, \theta_{1,p})$ (respectively $(\sigma_{2,p}, \theta_{2,p})$) is a biderivation of \mathfrak{n} (respectively \mathfrak{q}) follows directly from the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} . The identities $\mu\theta_{1,p} = \theta_{2,p}\mu$ and $\mu\sigma_{1,p} = \sigma_{2,p}\mu$ are immediate consequences of (LbEQ1) and (LbEQ2) respectively.

Observe that the combinations of the identities (LbCOM1) and (LbCOM4) and the identities (LbCOM5) and (LbCOM6) yield the equalities

$$-n^{(q^p)} = n^{(p^q)} \quad \text{and} \quad -{}^q(n^p) = {}^q(p^n).$$

These together with (LbCOM2)–(LbCOM5) allow us to prove that $((\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p}))$ does satisfy conditions (10)–(15) from Definition 3.5. Therefore, ψ is well defined, while it is obviously \mathbf{k} -linear. Moreover, due to (16) we know that

$$[\psi(p), \psi(p')] = ((\sigma_{1,p}\sigma_{1,p'} - \sigma_{1,p'}\sigma_{1,p}, \theta_{1,p}\sigma_{1,p'} - \sigma_{1,p'}\theta_{1,p}), (\sigma_{2,p}\sigma_{2,p'} - \sigma_{2,p'}\sigma_{2,p}, \theta_{2,p}\sigma_{2,p'} - \sigma_{2,p'}\theta_{2,p})),$$

and by definition

$$\psi([p, p']) = ((\sigma_{1,[p,p']}, \theta_{1,[p,p']}), (\sigma_{2,[p,p']}, \theta_{2,[p,p']})).$$

One can easily check that the corresponding components are equal by making use of the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} . Hence, ψ is a homomorphism of Leibniz algebras.

Recall that

$$\begin{aligned}\Delta\varphi(m) &= ((d_m\mu, D_m\mu), (\mu d_m, \mu D_m)), \\ \psi\eta(m) &= ((\sigma_{1,\eta(m)}, \theta_{1,\eta(m)}), (\sigma_{2,\eta(m)}, \theta_{2,\eta(m)})),\end{aligned}$$

for any $m \in \mathfrak{m}$, but

$$\begin{aligned}d_m\mu(n) &= -\xi_2(\mu(n), m) = -n^m = -n^{\eta(m)} = \sigma_{1,\eta(m)}(n), \\ D_m\mu(n) &= \xi_1(m, \mu(n)) = {}^m n = \eta(m)n = \theta_{1,\eta(m)}(n), \\ \mu d_m(q) &= -\mu\xi_2(q, m) = -q^m = -q^{\eta(m)} = \sigma_{2,\eta(m)}(q), \\ \mu D_m(q) &= \mu\xi_1(m, q) = {}^m q = \eta(m)q = \theta_{2,\eta(m)}(q),\end{aligned}$$

for all $n \in \mathfrak{n}$, $q \in \mathfrak{q}$, due to (LbM1a), (LbM1b), (LbM2a), (LbM2b). Therefore, $\Delta\varphi = \psi\eta$.

It only remains to check the behaviour of (φ, ψ) regarding the action of \mathfrak{p} on \mathfrak{m} . Let $m \in \mathfrak{m}$ and $p \in \mathfrak{p}$. Due to (17) and (18),

$$\begin{aligned}\psi^{(p)}\varphi(m) &= (\sigma_{1,p}d_m - d_m\sigma_{2,p}, \theta_{1,p}d_m - d_m\theta_{2,p}), \\ \varphi(m)^{\psi(p)} &= (d_m\sigma_{2,p} - \sigma_{1,p}d_m, D_m\sigma_{2,p} - \sigma_{1,p}D_m).\end{aligned}$$

On the other hand, by definition, we know that

$$\begin{aligned}\varphi({}^p m) &= (d_{{}^p m}, D_{{}^p m}), \\ \varphi(m^p) &= (d_{(m^p)}, D_{(m^p)}).\end{aligned}$$

Directly from (LbM3a), (LbM3b), (LbM3c) and (LbM3d) one can easily confirm that the required identities between components hold. Hence, we can finally ensure that (φ, ψ) is a homomorphism of Leibniz crossed modules.

Now let us show that it is necessary that at least one of the conditions (CON1)–(CON3) holds in order to prove the converse statement. Let us suppose that there is a homomorphism of crossed modules

$$\begin{array}{ccc} \mathfrak{m} & \xrightarrow{\eta} & \mathfrak{p} \\ \varphi \downarrow & & \downarrow \psi \\ \text{Bider}(\mathfrak{q}, \mathfrak{n}) & \xrightarrow{\Delta} & \text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu) \end{array} \quad (24)$$

Given $m \in \mathfrak{m}$ and $p \in \mathfrak{p}$, let us denote $\varphi(m)$ by (d_m, D_m) and $\psi(p)$ by $((\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p}))$, which satisfy conditions (4)–(6) from Definition 3.1 and conditions (8)–(15) from Definition 3.5 respectively. Also, due to the definition of Δ (see Proposition 3.8), the commutativity of (24) can be expressed by the identity

$$((d_m\mu, D_m\mu), (\mu d_m, \mu D_m)) = ((\sigma_{1,\eta(m)}, \theta_{1,\eta(m)}), (\sigma_{2,\eta(m)}, \theta_{2,\eta(m)})), \quad (25)$$

for all $m \in \mathfrak{m}$. It is possible to define four bilinear maps, for which we use the same notation used for actions, from $\mathfrak{p} \times \mathfrak{n}$ to \mathfrak{n} , $\mathfrak{n} \times \mathfrak{p}$ to \mathfrak{n} , $\mathfrak{p} \times \mathfrak{q}$ to \mathfrak{q} and $\mathfrak{q} \times \mathfrak{p}$ to \mathfrak{q} , given by

$$\begin{aligned} {}^p n &= \theta_{1,p}(n), & n^p &= -\sigma_{1,p}(n), \\ {}^p q &= \theta_{2,p}(q), & q^p &= -\sigma_{2,p}(q), \end{aligned}$$

for all $n \in \mathfrak{n}$, $p \in \mathfrak{p}$, $q \in \mathfrak{q}$. These maps define actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} . The first three identities for the action on \mathfrak{n} (respectively \mathfrak{q}) follow easily from the fact that $(\sigma_{1,p}, \theta_{1,p})$ (respectively $(\sigma_{2,p}, \theta_{2,p})$) is a biderivation of \mathfrak{n} (respectively \mathfrak{q}).

Since ψ is a Leibniz homomorphism, we get that

$$\begin{aligned} ((\sigma_{1,[p,p']}, \theta_{1,[p,p']}), (\sigma_{2,[p,p']}, \theta_{2,[p,p']})) &= ((\sigma_{1,p}\sigma_{1,p'} - \sigma_{1,p'}\sigma_{1,p}, \theta_{1,p}\sigma_{1,p'} - \sigma_{1,p'}\theta_{1,p}), \\ &(\sigma_{2,p}\sigma_{2,p'} - \sigma_{2,p'}\sigma_{2,p}, \theta_{2,p}\sigma_{2,p'} - \sigma_{2,p'}\theta_{2,p})). \end{aligned}$$

The identities between the first and the second (respectively the third and the fourth) components in those quadruples allow us to confirm the fourth and fifth identities for the action of \mathfrak{p} on \mathfrak{n} (respectively \mathfrak{q}).

As for the last condition for both actions, it is fairly straightforward to check that

$$\begin{aligned} [p,p']n - ({}^p n)^{p'} &= \theta_{1,p}\sigma_{1,p'}(n), \\ [p,p']q - ({}^p q)^{p'} &= \theta_{2,p}\sigma_{2,p'}(q), \end{aligned}$$

while

$$\begin{aligned} {}^p(p' n) &= \theta_{1,p}\theta_{1,p'}(n), \\ {}^p(p' q) &= \theta_{2,p}\theta_{2,p'}(q), \end{aligned}$$

for all $n \in \mathfrak{n}$, $p, p' \in \mathfrak{p}$, $q \in \mathfrak{q}$. However, if at least one of the conditions (CON1)–(CON3) holds, due to Lemma 4.2 (i), $\theta_{1,p}\sigma_{1,p'}(n) = \theta_{1,p}\theta_{1,p'}(n)$ and $\theta_{2,p}\sigma_{2,p'}(q) = \theta_{2,p}\theta_{2,p'}(q)$. Therefore, we can ensure that there are Leibniz actions of \mathfrak{p} on both \mathfrak{n} and \mathfrak{q} , which induce actions of \mathfrak{m} on \mathfrak{n} and \mathfrak{q} via η .

The reader might have noticed that a fourth possible condition on $(\mathfrak{n}, \mathfrak{q}, \mu)$ could have been considered in order to guarantee the existence of the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} from the existence of the homomorphism of Leibniz crossed modules (φ, ψ) . In fact, if $[\mathfrak{n}, \mathfrak{n}] = \mathfrak{n}$ and $\text{Ann}(\mathfrak{q}) = 0$, the problem with the last condition for the actions could have been solved in the same way. Nevertheless, this fourth condition does not guarantee that (ii) holds, as we will prove immediately below.

Regarding (LbEQ1) and (LbEQ2), they follow directly from (9) (observe that, by hypothesis, $(\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p})$ is a biderivation of $(\mathfrak{n}, \mathfrak{q}, \mu)$ for any $p \in \mathfrak{p}$). Similarly, (LbCOM1)–(LbCOM6) follow almost immediately from (10)–(15). Hence, (i) holds.

Concerning (ii), we can define $\xi_1(m, q) = D_m(q)$ and $\xi_2(q, m) = -d_m(q)$ for any $m \in \mathfrak{m}$, $q \in \mathfrak{q}$. In this way, ξ_1 and ξ_2 are clearly bilinear. (LbM1a), (LbM1b), (LbM2a) and (LbM2b) follow immediately from the identity (25) and the fact that the actions of \mathfrak{m} on \mathfrak{n} and \mathfrak{q} are induced by the actions of \mathfrak{p} via η .

Identities (LbM5a), (LbM5b) and (LbM5c) are a direct consequence of (4)–(6) (recall that, by hypothesis, (d_m, D_m) is a biderivation from \mathfrak{q} to \mathfrak{n} for any $m \in \mathfrak{m}$).

Since φ is a Leibniz homomorphism, we have that

$$(d_{[m,m']}, D_{[m,m']}) = (d_m \mu d_{m'} - d_{m'} \mu d_m, D_m \mu d_{m'} - d_{m'} \mu D_m).$$

This identity, together with (LbM2a) and (LbM2b), allows to easily prove that (LbM4a) and (LbM4b) hold.

Note that, since (φ, ψ) is a homomorphism of Leibniz crossed modules, $\varphi({}^p m) = \psi({}^p \varphi(m))$ and $\varphi(m^p) = \varphi(m)^{\psi(p)}$ for all $m \in \mathfrak{m}$, $p \in \mathfrak{p}$. Due to the definition of the action of $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ on $\text{Bider}(\mathfrak{q}, \mathfrak{n})$ (see Theorem 3.9), we can write

$$\begin{aligned} (d_{({}^p m)}, D_{({}^p m)}) &= (\sigma_{1,p} d_m - d_m \sigma_{2,p}, \theta_{1,p} d_m - d_m \theta_{2,p}), \\ (d_{(m^p)}, D_{(m^p)}) &= (d_m \sigma_{2,p} - \sigma_{1,p} d_m, D_m \sigma_{2,p} - \sigma_{1,p} D_m). \end{aligned}$$

Identities (LbM3a), (LbM3b), (LbM3c) and (LbM3d) follow immediately from the previous identities.

Regarding (LbM6a) and (LbM6b), directly from the definition of ξ_1 , ξ_2 and the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} , we have that

$$\begin{aligned} \xi_1(m, {}^p q) &= D_m \theta_{2,p}(q), & {}^p \xi_1(m, q) &= \theta_{1,p} D_m(q), \\ -\xi_1(m, q^p) &= D_m \sigma_{2,p}(q), & -{}^p \xi_2(q, m) &= \theta_{1,p} d_m(q), \end{aligned}$$

for all $m \in \mathfrak{m}$, $p \in \mathfrak{p}$, $q \in \mathfrak{q}$. Nevertheless, if at least one of the conditions (CON1)–(CON3) holds, due to Lemma 4.2 (ii), $D_m \theta_{2,p}(q) = D_m \sigma_{2,p}(q)$ and $\theta_{1,p} D_m(q) = \theta_{1,p} d_m(q)$. Hence, (ii) holds. \blacksquare

4.4. REMARK. A closer look at the proof of the previous theorem shows that neither conditions (LbM6a) and (LbM6b), nor the identities ${}^p({}^{p'} n) = [{}^{p,p'}]n - ({}^p n)^{p'}$ and ${}^p({}^{p'} q) = [{}^{p,p'}]q - ({}^p q)^{p'}$ (which correspond to the sixth axiom satisfied by the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} respectively) are necessary in order to prove the existence of a homomorphism of crossed modules (φ, ψ) from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$, under the hypothesis that (i) and (ii) hold. Actually, if we remove those conditions from (i) and (ii), the converse statement would be true for any Leibniz crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$, even if it does not satisfy any of the conditions (CON1)–(CON3). The problem is that (LbM6a) and (LbM6b), together with the sixth identity satisfied by the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{q} are essential in order to prove that (i) and (ii) as in Theorem 4.3 describe a set of actions of $(\mathfrak{m}, \mathfrak{p}, \eta)$ on $(\mathfrak{n}, \mathfrak{q}, \mu)$, as we will show immediately below. This agrees with the idea of $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ not being “good enough” to be the actor of $(\mathfrak{n}, \mathfrak{q}, \mu)$ in general, just as $\text{Bider}(\mathfrak{m})$ is not always the actor of a Leibniz algebra \mathfrak{m} .

4.5. EXAMPLE. Let $(\mathfrak{m}, \mathfrak{p}, \eta) \in \mathbf{XLb}$, there is a homomorphism

$$(\varphi, \psi): (\mathfrak{m}, \mathfrak{p}, \eta) \rightarrow \overline{\text{Act}}(\mathfrak{m}, \mathfrak{p}, \eta)$$

, with $\varphi(m) = (d_m, D_m)$ and $\psi(p) = ((\sigma_{1,p}, \theta_{1,p}), (\sigma_{2,p}, \theta_{2,p}))$, where

$$d_m(p) = -{}^p m, \quad D_m(p) = m^p,$$

and

$$\begin{aligned} \sigma_{1,p}(m) &= -m^p, & \theta_{1,p}(m) &= {}^p m, \\ \sigma_{2,p}(p') &= -[p', p], & \theta_{2,p}(p') &= [p, p'], \end{aligned}$$

for all $m \in \mathfrak{m}$, $p, p' \in \mathfrak{p}$. Calculations in order to prove that (φ, ψ) is indeed a homomorphism of Leibniz crossed modules are fairly straightforward. Of course, this homomorphism does not necessarily define a set of actions from which it is possible to construct the semidirect product. Theorem 4.3, along with the result immediately bellow, shows that if $(\mathfrak{m}, \mathfrak{p}, \eta)$ satisfies at least one of the conditions (CON1)–(CON3), then the previous homomorphism does define an appropriate set of actions of $(\mathfrak{m}, \mathfrak{p}, \eta)$ on itself.

Let $(\mathfrak{m}, \mathfrak{p}, \eta)$ and $(\mathfrak{n}, \mathfrak{q}, \mu)$ be Leibniz crossed modules such that (i) and (ii) from Theorem 4.3 hold. Therefore, there are Leibniz actions of \mathfrak{m} on \mathfrak{n} and of \mathfrak{p} on \mathfrak{q} , so it makes sense to consider the semidirect products of Leibniz algebras $\mathfrak{n} \rtimes \mathfrak{m}$ and $\mathfrak{q} \rtimes \mathfrak{p}$. Furthermore, we have the following result.

4.6. THEOREM. *There is an action of the Leibniz algebra $\mathfrak{q} \rtimes \mathfrak{p}$ on the Leibniz algebra $\mathfrak{n} \rtimes \mathfrak{m}$, given by*

$${}^{(q,p)}(n, m) = ({}^q n + {}^p n + \xi_2(q, m), {}^p m), \quad (26)$$

$$(n, m)^{(q,p)} = (n^q + n^p + \xi_1(m, q), m^p), \quad (27)$$

for all $(q, p) \in \mathfrak{q} \rtimes \mathfrak{p}$, $(n, m) \in \mathfrak{n} \rtimes \mathfrak{m}$, with ξ_1 and ξ_2 as in Theorem 4.3. Moreover, the Leibniz homomorphism $(\mu, \eta): \mathfrak{n} \rtimes \mathfrak{m} \rightarrow \mathfrak{q} \rtimes \mathfrak{p}$, given by

$$(\mu, \eta)(n, m) = (\mu(n), \eta(m)),$$

for all $(n, m) \in \mathfrak{n} \rtimes \mathfrak{m}$, together with the previous action, is a Leibniz crossed module.

PROOF. Identities (26) and (27) follow easily from the conditions satisfied by $(\mathfrak{m}, \mathfrak{p}, \eta)$ and $(\mathfrak{n}, \mathfrak{q}, \mu)$ (see Theorem 4.3). Nevertheless, as an example, we show how to prove the third one. Calculations for the rest of the identities are similar. Let $(n, m), (n', m') \in \mathfrak{n} \rtimes \mathfrak{m}$ and $(q, p) \in \mathfrak{q} \rtimes \mathfrak{p}$. By routine calculations we get that

$$\begin{aligned} [(n, m), (n', m')]^{(q,p)} &= \underbrace{[n, n'^q]}_{(1)} + \underbrace{[n, n'^p]}_{(2)} + \underbrace{[n, \xi_1(m', q)]}_{(3)} + \underbrace{m(n'^q)}_{(4)} \\ &\quad + \underbrace{m(n'^p)}_{(5)} + \underbrace{m\xi_1(m', q)}_{(6)} + \underbrace{n^{(m'p)}}_{(7)} + \underbrace{[m, m'^p]}_{(8)}, \end{aligned}$$

$$[(n, m), (n', m')]^{(q,p)} = \underbrace{[n, n']^q}_{(1')} + \underbrace{[n, n']^p}_{(2')} + \underbrace{(n^{m'})^q}_{(3')} + \underbrace{({}^m n')^q}_{(4')} \\ + \underbrace{({}^m n')^p}_{(5')} + \underbrace{\xi_1([m, m'], q)}_{(6')} + \underbrace{(n^{m'})^p}_{(7')} + \underbrace{[m, m']^p}_{(8')},$$

$$[(n, m)^{(q,p)}, (n', m')] = \underbrace{[n^q, n']}_{(1'')} + \underbrace{[n^p, n']}_{(2'')} + \underbrace{(n^q)^{m'}}_{(3'')} + \underbrace{[\xi_1(m, q), n']}_{(4'')} \\ + \underbrace{({}^{m^p})n'}_{(5'')} + \underbrace{\xi_1(m, q)^{m'}}_{(6'')} + \underbrace{(n^p)^{m'}}_{(7'')} + \underbrace{[m^p, m']}_{(8'')}.$$

Let us show that $(i) = (i') - (i'')$ for $i = 1, \dots, 8$. It is immediate for $i = 1, 2, 8$ due to the action of \mathfrak{q} on \mathfrak{n} and the actions of \mathfrak{p} on \mathfrak{n} and \mathfrak{m} . For $i = 5$, the identity follows from the fact that the action of \mathfrak{m} on \mathfrak{n} is defined via η together with the equivariance of η . Namely,

$${}^m(n^p) = \eta^{(m)}(n^p) = (\eta^{(m)}n^p) - [{}^{\eta(m), p}n^p] = ({}^m n^p) - \eta^{(m^p)}n^p = ({}^m n^p) - ({}^{m^p})n^p.$$

The procedure is similar for $i = 7$. For $i = 3$, it is necessary to make use of the Peiffer identity of μ , (LbM1b), the definition of the action of \mathfrak{m} on \mathfrak{n} and \mathfrak{q} via η and (LbCOM1):

$$[n, \xi_1(m', q)] = n^{\mu \xi_1(m', q)} = n^{(m' q)} = n^{(\eta(m') q)} = (n^{\eta(m')})^q - (n^q)^{\eta(m')} = (n^{m'})^q - (n^{m'})^q.$$

The conditions required in order to prove the identity for $i = 4$ are the same used for $i = 3$ except (LbCOM1), which is replaced by (LbCOM2).

Finally, for $i = 6$, due to (LbM4b) and the definition of the action of \mathfrak{m} on \mathfrak{n} via η , we know that

$$\xi_1([m, m'], q) = \xi_1(m, q)^{m'} - {}^m \xi_2(q, m') = \xi_1(m, q)^{m'} - \eta^{(m)} \xi_2(q, m'),$$

but applying (LbM6b), we get

$$\xi_1([m, m'], q) = \xi_1(m, q)^{m'} + \eta^{(m)} \xi_1(m', q) = \xi_1(m, q)^{m'} + {}^m \xi_1(m', q),$$

so $(6) = (6') - (6'')$ and the third identity holds. Note that (LbM6a) and (LbM6b) are necessary in order to check the fourth and fifth identities respectively.

Checking that (μ, η) is indeed a Leibniz homomorphism follows directly from the definition of the action of \mathfrak{m} on \mathfrak{n} via η together with the conditions (LbEQ1) and (LbEQ2). Regarding the equivariance of (μ, η) , given $(n, m) \in \mathfrak{n} \rtimes \mathfrak{m}$ and $(q, p) \in \mathfrak{q} \rtimes \mathfrak{p}$,

$$\begin{aligned} (\mu, \eta)^{(q,p)}(n, m) &= (\mu, \eta)^{(q,n + p n + \xi_2(q, m), p m)} \\ &= (\mu^{(q,n)} + \mu^{(p,n)} + \mu \xi_2(q, m), \eta^{(p,m)}) \\ &= ([q, \mu(n)] + {}^p \mu(n) + q^m, [p, \eta(m)]) \\ &= ([q, \mu(n)] + {}^p \mu(n) + q^{\eta(m)}, [p, \eta(m)]) \\ &= [(q, p), (\mu(n), \eta(m))], \end{aligned}$$

due to the equivariance of μ and η , (LbEQ1), (LbM1a) and the definition of the action of \mathfrak{m} on \mathfrak{q} via η . Similarly, but using (LbEQ2) and (LbM1b) instead of (LbEQ1) and (LbM1a), it can be proved that $(\mu, \eta)((n, m)^{(q,p)}) = [(\mu(n), \eta(m)), (q, p)]$.

The Peiffer identity of (μ, η) follows easily from the homonymous property of μ and η , the definition of the action of \mathfrak{m} on \mathfrak{n} via η and the conditions (LbM2a) and (LbM2b). ■

4.7. DEFINITION. *The Leibniz crossed module $(\mathfrak{n} \rtimes \mathfrak{m}, \mathfrak{q} \rtimes \mathfrak{p}, (\mu, \eta))$ is called the semidirect product of the Leibniz crossed modules $(\mathfrak{n}, \mathfrak{q}, \mu)$ and $(\mathfrak{m}, \mathfrak{p}, \eta)$.*

Note that the semidirect product determines an obvious split extension of $(\mathfrak{m}, \mathfrak{p}, \eta)$ by $(\mathfrak{n}, \mathfrak{q}, \mu)$

$$(0, 0, 0) \longrightarrow (\mathfrak{n}, \mathfrak{q}, \mu) \longrightarrow (\mathfrak{n} \rtimes \mathfrak{m}, \mathfrak{q} \rtimes \mathfrak{p}, (\mu, \eta)) \begin{array}{c} \longleftarrow \\ \longrightarrow \end{array} (\mathfrak{m}, \mathfrak{p}, \eta) \longrightarrow (0, 0, 0)$$

Conversely, any split extension of $(\mathfrak{m}, \mathfrak{p}, \eta)$ by $(\mathfrak{n}, \mathfrak{q}, \mu)$ is isomorphic to their semidirect product, where the action of $(\mathfrak{m}, \mathfrak{p}, \eta)$ on $(\mathfrak{n}, \mathfrak{q}, \mu)$ is induced by the splitting homomorphism.

4.8. REMARK. If $(\mathfrak{m}, \mathfrak{p}, \eta)$ and $(\mathfrak{n}, \mathfrak{q}, \mu)$ are Leibniz crossed modules and at least one of the following conditions holds,

1. $\text{Ann}(\mathfrak{n}) = 0 = \text{Ann}(\mathfrak{q})$,
2. $\text{Ann}(\mathfrak{n}) = 0$ and $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$,
3. $[\mathfrak{n}, \mathfrak{n}] = \mathfrak{n}$ and $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$,

an action of the crossed module $(\mathfrak{m}, \mathfrak{p}, \eta)$ on $(\mathfrak{n}, \mathfrak{q}, \mu)$ can be also defined as a homomorphism of Leibniz crossed modules from $(\mathfrak{m}, \mathfrak{p}, \eta)$ to $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$. In other words, under one of those conditions, $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ is the actor of $(\mathfrak{n}, \mathfrak{q}, \mu)$ and it can be denoted simply by $\text{Act}(\mathfrak{n}, \mathfrak{q}, \mu)$.

4.9. EXAMPLE.

(i) Let \mathfrak{n} be an ideal of a Leibniz algebra \mathfrak{q} and consider the crossed module $(\mathfrak{n}, \mathfrak{q}, \iota)$, where ι is the inclusion. It is easy to check that $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \iota) = (X, Y, \iota)$, where X is a Leibniz algebra isomorphic to $\{(d, D) \in \text{Bider}(\mathfrak{q}) \mid d(q), D(q) \in \mathfrak{n} \text{ for all } q \in \mathfrak{q}\}$ and Y is a Leibniz algebra isomorphic to $\{(d, D) \in \text{Bider}(\mathfrak{q}) \mid (d|_{\mathfrak{n}}, D|_{\mathfrak{n}}) \in \text{Bider}(\mathfrak{n})\}$.

(ii) Given a Leibniz algebra \mathfrak{q} , it can be regarded as a Leibniz crossed module in two obvious ways, $(0, \mathfrak{q}, 0)$ and $(\mathfrak{q}, \mathfrak{q}, \text{id}_{\mathfrak{q}})$. As a particular case of the previous example, one can easily check that $\overline{\text{Act}}(0, \mathfrak{q}, 0) \cong (0, \text{Bider}(\mathfrak{q}), 0)$ and $\overline{\text{Act}}(\mathfrak{q}, \mathfrak{q}, \text{id}_{\mathfrak{q}}) \cong (\text{Bider}(\mathfrak{q}), \text{Bider}(\mathfrak{q}), \text{id})$.

(iii) An action of a Leibniz crossed module on an abelian Leibniz crossed module is precisely a Leibniz crossed module representation ([7])

(iv) Every Lie crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$ can be regarded as a Leibniz crossed module (see for instance [9, Remark 3.9]). Note that in this situation, both the multiplication and the action are antisymmetric. The actor of $(\mathfrak{n}, \mathfrak{q}, \mu)$ is $(\text{Der}(\mathfrak{q}, \mathfrak{n}), \text{Der}(\mathfrak{n}, \mathfrak{q}, \mu), \Delta)$, where $\text{Der}(\mathfrak{q}, \mathfrak{n})$ is the Lie algebra of all derivations from \mathfrak{q} to \mathfrak{n} and $\text{Der}(\mathfrak{n}, \mathfrak{q}, \mu)$ is the Lie

algebra of derivations of the crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$ (see [10] for the details). Given $(d, D) \in \text{Bider}(\mathfrak{q}, \mathfrak{n})$, both d and D are elements in $\text{Der}(\mathfrak{q}, \mathfrak{n})$. Additionally, if we assume that at least one of the conditions from the previous lemma holds, then either $\text{Ann}(\mathfrak{n}) = 0$ or $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{q}$. In this situation, one can easily derive from (6) that $\text{Bider}(\mathfrak{q}, \mathfrak{n}) = \{(d, d) \mid d \in \text{Der}(\mathfrak{q}, \mathfrak{n})\}$. Besides, the bracket in $\text{Bider}(\mathfrak{q}, \mathfrak{n})$ becomes antisymmetric and, as a Lie algebra, it is isomorphic to $\text{Der}(\mathfrak{q}, \mathfrak{n})$. Similarly, $\text{Bider}(\mathfrak{n}, \mathfrak{q}, \mu)$ is a Lie algebra isomorphic to $\text{Der}(\mathfrak{n}, \mathfrak{q}, \mu)$ and $\overline{\text{Act}}(\mathfrak{n}, \mathfrak{q}, \mu)$ is a Lie crossed module isomorphic to $\text{Act}(\mathfrak{n}, \mathfrak{q}, \mu)$.

5. Center of a Leibniz crossed module

Let us assume in this section that $(\mathfrak{n}, \mathfrak{q}, \mu)$ is a Leibniz crossed module that satisfies at least one of the conditions (CON1)–(CON3). Denote by $Z(\mathfrak{q})$ the center of the Leibniz algebra \mathfrak{q} , which in this case coincides with its annihilator (note that the center and the annihilator are not the same object in general). Consider the canonical homomorphism (φ, ψ) from $(\mathfrak{n}, \mathfrak{q}, \mu)$ to $\text{Act}(\mathfrak{n}, \mathfrak{q}, \mu)$, as in Example 4.5. It is easy to check that

$$\text{Ker}(\varphi) = \mathfrak{n}^{\mathfrak{q}} \quad \text{and} \quad \text{Ker}(\psi) = \text{st}_{\mathfrak{q}}(\mathfrak{n}) \cap Z(\mathfrak{q}),$$

where $\mathfrak{n}^{\mathfrak{q}} = \{n \in \mathfrak{n} \mid {}^q n = n^{\mathfrak{q}} = 0, \text{ for all } q \in \mathfrak{q}\}$ and $\text{st}_{\mathfrak{q}}(\mathfrak{n}) = \{q \in \mathfrak{q} \mid {}^q n = n^{\mathfrak{q}} = 0, \text{ for all } n \in \mathfrak{n}\}$. Therefore, the kernel of (φ, ψ) is the Leibniz crossed module $(\mathfrak{n}^{\mathfrak{q}}, \text{st}_{\mathfrak{q}}(\mathfrak{n}) \cap Z(\mathfrak{q}), \mu)$. Thus, the kernel of (φ, ψ) coincides with the center of the crossed module $(\mathfrak{n}, \mathfrak{q}, \mu)$, as defined in the preliminary version of [1, Definition 27] for crossed modules in modified categories of interest. This definition of center agrees with the categorical notion of center by Huq [11] and confirms that our construction of the actor for a Leibniz crossed module is consistent.

5.1. EXAMPLE. Consider the crossed module $(\mathfrak{n}, \mathfrak{q}, \iota)$, where \mathfrak{n} is an ideal of \mathfrak{q} and ι is the inclusion. Then, its center is given by the Leibniz crossed module $(\mathfrak{n} \cap Z(\mathfrak{q}), Z(\mathfrak{q}), \iota)$. In particular, the center of $(0, \mathfrak{q}, 0)$ is $(0, Z(\mathfrak{q}), 0)$ and the center of $(\mathfrak{q}, \mathfrak{q}, \text{id}_{\mathfrak{q}})$ is $(Z(\mathfrak{q}), Z(\mathfrak{q}), \text{id})$.

By analogy to the definitions given for crossed modules of Lie algebras (see [10]), we can define the crossed module of *inner biderivations* of $(\mathfrak{n}, \mathfrak{q}, \mu)$, denoted by $\text{InnBider}(\mathfrak{n}, \mathfrak{q}, \mu)$, as $\text{Im}(\varphi, \psi)$, which is obviously an ideal. The crossed module of *outer biderivations*, denoted by $\text{OutBider}(\mathfrak{n}, \mathfrak{q}, \mu)$, is the quotient of $\text{Act}(\mathfrak{n}, \mathfrak{q}, \mu)$ by $\text{InnBider}(\mathfrak{n}, \mathfrak{q}, \mu)$.

Let

$$(0, 0, 0) \longrightarrow (\mathfrak{n}, \mathfrak{q}, \mu) \longrightarrow (\mathfrak{n}', \mathfrak{q}', \mu') \longrightarrow (\mathfrak{n}'', \mathfrak{q}'', \mu'') \longrightarrow (0, 0, 0)$$

be a short exact sequence of crossed modules of Leibniz algebras. Then, there exists a homomorphism of Leibniz crossed modules $(\alpha, \beta): (\mathfrak{n}', \mathfrak{q}', \mu') \rightarrow \text{Act}(\mathfrak{n}, \mathfrak{q}, \mu)$ so that the following diagram is commutative:

$$\begin{array}{ccccccccc} (0, 0, 0) & \longrightarrow & (\mathfrak{n}, \mathfrak{q}, \mu) & \longrightarrow & (\mathfrak{n}', \mathfrak{q}', \mu') & \longrightarrow & (\mathfrak{n}'', \mathfrak{q}'', \mu'') & \longrightarrow & (0, 0, 0) \\ & & \downarrow & & \downarrow (\alpha, \beta) & & \downarrow & & \\ (0, 0, 0) & \longrightarrow & \text{InnBider}(\mathfrak{n}, \mathfrak{q}, \mu) & \longrightarrow & \text{Act}(\mathfrak{n}, \mathfrak{q}, \mu) & \longrightarrow & \text{OutBider}(\mathfrak{n}, \mathfrak{q}, \mu) & \longrightarrow & (0, 0, 0) \end{array}$$

where (α, β) is defined as $\alpha(n') = (d_{n'}, D_{n'})$ and $\beta(q') = ((\sigma_{1,q'}, \theta_{1,q'}), (\sigma_{2,q'}, \theta_{2,q'}))$, with

$$d_{n'}(q) = -{}^q n', \quad D_{n'}(q) = n'^q,$$

and

$$\begin{aligned} \sigma_{1,q'}(n) &= -n^{q'}, & \theta_{1,q'}(n) &= {}^{q'} n, \\ \sigma_{2,q'}(q) &= -[q, q'], & \theta_{2,q'}(q) &= [q', q], \end{aligned}$$

for all $n' \in \mathfrak{n}'$, $q' \in \mathfrak{q}'$, $n \in \mathfrak{n}$, $q \in \mathfrak{q}$.

References

- [1] Uslu, E Ö and Çetin, S and Arslan, A F. On crossed modules in modified categories of interest. *Math. Commun.*, 2017, 22(1): 103–119
- [2] Borceux F, Janelidze G, Kelly G M. On the representability of actions in a semi-abelian category. *Theory Appl Categ*, 2005, 14: 244–286
- [3] Borceux F, Janelidze G, Kelly G M. Internal object actions. *Comment. Math. Univ. Carolin.*, 2005, 46(2): 235–255
- [4] Boyaci Y, Casas J M, Datuashvili T, Uslu E Ö. Actions in modified categories of interest with application to crossed modules. *Theory Appl. Categ.*, 2015, 30: 882–908
- [5] Casas J M, Casado R F, Khmaladze E, Ladra M. Universal enveloping crossed module of a Lie crossed module. *Homology Homotopy Appl.*, 2014, 16(2): 143–158
- [6] Casas J M, Datuashvili T, Ladra M. Universal strict general actors and actors in categories of interest. *Appl. Categ. Structures*, 2010, 18(1): 85–114
- [7] Fernández-Casado R, García-Martínez, X, Ladra M. A Natural Extension of the Universal Enveloping Algebra Functor to Crossed Modules of Leibniz Algebras. *Appl. Categ. Structures*, 2017. doi:10.1007/s10485-016-9472-9
- [8] Casas J M, Inassaridze N, Khmaladze E, Ladra M. Adjunction between crossed modules of groups and algebras. *J. Homotopy Relat. Struct.*, 2014, 9(1): 223–237
- [9] Casas J M, Khmaladze E, Ladra M. Low-dimensional non-abelian Leibniz cohomology. *Forum Math.*, 2013, 25: 443–469
- [10] Casas J M, Ladra M. The actor of a crossed module in Lie algebras. *Comm. Algebra*, 1998, 26(7): 2065–2089
- [11] Huq S A. Commutator, nilpotency, and solvability in categories. *Quart. J. Math. Oxford Ser. (2)*, 1968, 19: 363–389

- [12] Loday J L. Une version non commutative des algèbres de Lie: les algèbres de Leibniz. Enseign. Math. (2), 1993, 39(3-4): 269–293
- [13] Loday J L, Pirashvili T. Universal enveloping algebras of Leibniz algebras and (co)homology. Math. Ann., 1993, 296(1): 139–158
- [14] Montoli A. Action accessibility for categories of interest. Theory Appl. Categ., 2010, 23(1): 7–21
- [15] Norrie K. Actions and automorphisms of crossed modules. Bull. Soc. Math. France, 1990, 118(2): 129–146
- [16] Orzech G. Obstruction theory in algebraic categories. I, II. J. Pure Appl. Algebra, 1972, 2: 287–314; 1972, 2: 315–340

*Department of Applied Mathematics I, University of Vigo
E. E. Forestal, 36005 Pontevedra, Spain*

*Department of Mathematics, University of Santiago de Compostela
Lope Gómez de Marzoa s/n 15782, Santiago de Compostela, Spain*

*Department of Mathematics, University of Santiago de Compostela
Lope Gómez de Marzoa s/n 15782, Santiago de Compostela, Spain*

*A. Razmadze Mathematical Institute, Tbilisi State University
0177 Tbilisi, Georgia*

Email: `jmcasas@uvigo.es`
`rapha.fdez@gmail.com`
`xabier.garcia@usc.es`
`e.khmal@gmail.com`

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available from the journal's server at <http://www.tac.mta.ca/tac/>. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$ is required. Articles in PDF format may be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ NICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca